



## Understanding the growth kinetics of *Listeria monocytogenes* on pressurized cooked meat products depending on its physiological state

Hereu, A.; Dalgaard, Paw; Garriga, M.; Aymerich, T.; Bover-Cid, S.

*Published in:*

2nd Workshop on Food Safety Technologies and Innovations Applied to Food Safety

*Publication date:*

2012

*Document Version*

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Hereu, A., Dalgaard, P., Garriga, M., Aymerich, T., & Bover-Cid, S. (2012). Understanding the growth kinetics of *Listeria monocytogenes* on pressurized cooked meat products depending on its physiological state. In *2nd Workshop on Food Safety Technologies and Innovations Applied to Food Safety: Book of Abstracts* (pp. 32). SICURA.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## UNDERSTANDING THE GROWTH KINETICS OF *LISTERIA MONOCYTOGENES* ON PRESSURIZED COOKED MEAT PRODUCTS DEPENDING ON ITS PHYSIOLOGICAL STATE

**A. Hereu<sup>1</sup>, P. Dalgaard<sup>2</sup>, M. Garriga<sup>1</sup>, T. Aymerich<sup>1</sup>, S. Bover-Cid<sup>1</sup>**

<sup>1</sup>IRTA. Food Safety Programme. Finca Camps i Armet, E-17121 Monells, Spain. <sup>2</sup>Technical University of Denmark (DTU), National Food Institute, Soltofts Plads, Bygning 221, DK-2800, Kgs. Lyngby, Denmark.

**Abstract.** The bactericidal effect of high hydrostatic pressure (HHP) on ready-to-eat (RTE) products has been largely studied. However, most available predictive models deal with HHP-inactivation, without considering the recovery of surviving cells during subsequent chilled storage, neither the influence of the physiological state of the pathogen. Therefore, we aimed to model the growth kinetics of *L. monocytogenes* on pressurized RTE sliced cooked meat products (cooked ham and mortadella), depending on the inoculum level and its physiological state, established by using different pre-culture conditions. Slices of both products were inoculated with *L. monocytogenes* CTC1034, at  $10^7$  and  $10^4$  cfu/g using two different pre-cultures in the stationary phase: one was adapted to refrigeration at 8°C and the other was cold shocked at -80 °C. Inoculated samples were vacuum packaged, pressurized (400 MPa, 5 min, 15°C) and thereafter stored at 4, 8 and 12 °C. *L. monocytogenes* was periodically enumerated on ALOA plates. Log-transformed counts were fitted to the Logistic model with delay to estimate the lag phase ( $\lambda$ ), maximum specific growth rate ( $\mu_{max}$ ) and maximum population density ( $N_{max}$ ). The influence of the storage temperature on  $\lambda$ ,  $\mu_{max}$  and  $N_{max}$  was assessed by secondary modeling. HHP-inactivation was influenced by product characteristics and the physiological state of the pathogen. Cold shock resulted in a significant baroprotection. Regardless of the inoculum level and the physiological state of *L. monocytogenes*, HHP had no effect on  $\mu_{max}$ . However a significant  $\lambda$  was observed for the pathogen when previously shocked at -80 °C. The developed secondary models for  $\mu_{max}$  and  $\lambda$ , allowed growth of *L. monocytogenes* during the storage of pressurized RTE cooked meat products to be predicted. In conclusion, the relevance of the physiological state of *L. monocytogenes* prior to HHP treatments is an important factor to be taken into account when assessing the effectiveness of HHP as listericidal post-processing treatment.